


Claims:

1. A compound of formula (I)



in which

- ▶ n represents 1, 2 or 3;
- ▶ A represents a substituent chosen from -C(O)-, -C(S)-, -CH₂-, -CHR¹⁰-, -CR¹⁰R¹¹-, -C(O)O-, -C(O)S-, -C(S)O-, -C(S)S-, -C(O)NH-, -C(NH)NH- and -C(S)NH-;
- ▶ B represents
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a naphthylene;
 - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
 - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a biphenylene;
 - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R¹² and R¹³ chosen, independently of each other, from halogen, CN, C(O)OR¹⁴, C(O)NR¹⁵R¹⁶, CF₃, OCF₃, -NO₂, N₃, OR¹⁴, SR¹⁴, NR¹⁵R¹⁶ and C₁₋₆-alkyl;

- ▶ C represents a substituent chosen from -O-, -S-, -CH₂-, -CHR¹⁷-, -CR¹⁷R¹⁸-, -NH- and -NR¹⁹;
- ▶ D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;
- ▶ E and G represent, independently of each other, a substituent chosen from H, OH, OR²⁰, NH₂ and NHR²⁰;

- R¹ represents a substituent chosen from H, C₁₋₆-alkyl, C(O)H and C(O)CH₃;
- R², R³, R⁶, R¹⁴, R¹⁵, R¹⁶ and R¹⁹ represent, independently of each other, a substituent chosen from H, C₁₋₆-alkyl, C(O)C₁₋₆-alkyl, -C(S)C₁₋₆-alkyl, -C(O)OC₁₋₆-alkyl, -C(O)NH₂, -C(S)NH₂, -C(NH)NH₂, -C(O)NHC₁₋₆-alkyl, -C(S)NHC₁₋₆-alkyl and -C(NH)NHC₁₋₆-alkyl;
- 5 ► R⁴ represents a substituent chosen from H, C₁₋₆-alkyl and R²¹;
- R⁵ represents a substituent chosen from H, C₁₋₆-alkyl, fucosyl and R²²;
- R⁷ represents a substituent chosen from H, C₁₋₆-alkyl, arabinosyl and R²³;
- R⁸ represents a substituent chosen from H, C₁₋₆-alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄ and R²⁴;
- 10 ► R⁹ represents a substituent chosen from H, C₁₋₆-alkyl, mannose, glycerol and R²⁵;
- R¹⁰, R¹¹, R¹⁷ and R¹⁸ represent, independently of each other, a substituent chosen from C₁₋₆-alkyl and F;
- R²⁰, R²¹, R²², R²³, R²⁴ and R²⁵ represent, independently of each other, a substituent chosen from C(O)C₁₋₆-alkyl, -C(S)C₁₋₆-alkyl, -C(O)OC₁₋₆-alkyl, -C(O)NH₂, -C(S)NH₂, -C(NH)NH₂, -C(O)NHC₁₋₆-alkyl, -C(S)NHC₁₋₆-alkyl and -C(NH)NHC₁₋₆-alkyl;
- 15

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are agriculturally acceptable, such as lithium, sodium, potassium and tetraalkylammonium salts.

2. The compound of formula (I) as claimed in claim 1, having one or other of the following characteristics, taken separately or in combination:

- n represents 2 or 3;
- A represents -C(O)- or -CH₂-;
- B represents a phenylene;
- 25 ► C represents -O-;
- D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- E and G represent NHC(O)CH₃;
- R¹ represents H, CH₃ or C(O)CH₃;
- 30 ► R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

3. The compound of formula (I) as claimed in claim 1 or 2, simultaneously having the following characteristics:

- n represents 2 or 3;

- ▶ A represents -C(O)- or -CH₂-;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

4. The compound of formula (I) as claimed in any one of claims 1 to 3, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

5. The compound of formula (I) as claimed in any one of claims 1 to 4, simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

6. The compound of formula (I) as claimed in any one of claims 1 to 5, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;

- B represents a phenylene;
- C represents -O-;
- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent NHC(O)CH_3 ;
- R^1 represents H, CH_3 or C(O)CH_3 ;
- R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

7. The compound as claimed in claim 1 and of formula (Ia)



(Ia)

in which

- n represents 1, 2 or 3,
- B represents
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a naphthylene;
 - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
 - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a biphenylene;
 - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, C(O)OR^{14} , $\text{C(O)NR}^{15}\text{R}^{16}$, CF_3 , OCF_3 , $-\text{NO}_2$, N_3 , OR^{14} , SR^{14} , $\text{NR}^{15}\text{R}^{16}$ and $\text{C}_{1-6}\text{-alkyl}$;

- C represents a substituent chosen from -O-, -S-, -CH₂-, -CHR¹⁷-, -CR¹⁷R¹⁸-, -NH- or -NR¹⁹;
- D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;
- E and G represent, independently of each other, a substituent chosen from H, OH, OR²⁰, NH₂, NHR²⁰;

- R¹ represents a substituent chosen from H, C₁₋₆-alkyl, C(O)H, and C(O)CH₃;

R², R³, and R⁶ represent, independently of each other, a substituent chosen from H, C₁₋₆-alkyl, C(O)C₁₋₆-alkyl, -C(S)C₁₋₆-alkyl, -C(O)OC₁₋₆-alkyl, -C(O)NH₂, -C(S)NH₂, -C(NH)NH₂, -C(O)NHC₁₋₆-alkyl, -C(S)NHC₁₋₆-alkyl or -C(NH)NHC₁₋₆-alkyl;

- R⁴ represents a substituent chosen from H, C₁₋₆-alkyl or R²¹;

- R⁵ represents a substituent chosen from H, C₁₋₆-alkyl, fucosyl or R²²;

- R⁷ represents a substituent chosen from H, C₁₋₆-alkyl, arabinosyl or R²³;

- R⁸ represents a substituent chosen from H, C₁₋₆-alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄ or R²⁴;

- R⁹ represents a substituent chosen from H, C₁₋₆-alkyl, mannose, glycerol or R²⁵;

- R¹⁰, R¹¹, R¹⁷ and R¹⁸ represent, independently of each other, a substituent chosen from C₁₋₆-alkyl or F;

- R¹⁴, R¹⁵, R¹⁶ and R¹⁹ represent, independently of each other, a substituent chosen from H or C₁₋₆-alkyl, -C(O)C₁₋₆-alkyl, -C(S)C₁₋₆-alkyl, -C(O)OC₁₋₆-alkyl, -C(O)NH₂, -C(S)NH₂, -C(NH)NH₂, -C(O)NHC₁₋₆-alkyl, -C(S)NHC₁₋₆-alkyl or -C(NH)NHC₁₋₆-alkyl;

- R²⁰, R²¹, R²², R²³, R²⁴ and R²⁵ represent, independently of each other, a substituent chosen from C(O)C₁₋₆-alkyl, -C(S)C₁₋₆-alkyl, -C(O)OC₁₋₆-alkyl, -C(O)NH₂, -C(S)NH₂, -C(NH)NH₂, -C(O)NHC₁₋₆-alkyl, -C(S)NHC₁₋₆-alkyl or -C(NH)NHC₁₋₆-alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

8. The compound of formula (Ia) as claimed in claim 7, having one or other of the following characteristics, taken separately or in combination:

- n represents 2 or 3;
- B represents a phenylene;
- C represents -O-;
- D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- E and G represent NHC(O)CH₃;

- ▶ R¹ represents H or CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

5

9. The compound of formula (Ia) as claimed in claim 7 or 8, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H or CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

10

10. The compound of formula (Ia) as claimed in any one of claims 7 to 9, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H or CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

20

11. The compound of formula (Ia) as claimed in any one of claims 7 to 10, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H or CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

25


30

35

12. The compound of formula (Ia) as claimed in any one of claims 7 to 11, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ C represents -O-;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or CH_3 ;
- ▶ R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

13. The compound as claimed in claim 1 and of formula (Ib)



(Ib)

in which

- ▶ n represents 1, 2 or 3,
- ▶ B represents
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a naphthylene;
 - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
 - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a biphenylene;

- or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl;

► C represents a substituent chosen from $-O-$, $-S-$, $-CH_2-$, $-CHR^{17}-$, $-CR^{17}R^{18}-$, $-NH-$ or $-NR^{19}$;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;

► E and G represent, independently of each other, a substituent chosen from H, OH, OR^{20} , NH_2 , NHR^{20} ;

► R^1 represents a substituent chosen from H, C_{1-6} -alkyl, $C(O)H$, and $C(O)CH_3$;

► R^2 , R^3 , and R^6 represent, independently of each other, a substituent chosen from H, C_{1-6} -alkyl, $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^4 represents a substituent chosen from H, C_{1-6} -alkyl or R^{21} ;

► R^5 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl or R^{22} ;

► R^7 represents a substituent chosen from H, C_{1-6} -alkyl, arabinosyl or R^{23} ;

► R^8 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$ or R^{24} ;

► R^9 represents a substituent chosen from H, C_{1-6} -alkyl, mannose, glycerol or R^{25} ;

► R^{10} , R^{11} , R^{17} and R^{18} represent, independently of each other, a substituent chosen from C_{1-6} -alkyl or F;

► R^{14} , R^{15} , R^{16} and R^{19} represent, independently of each other, a substituent chosen from H or C_{1-6} -alkyl, $-C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} represent, independently of each other, a substituent chosen from $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

14. The compound of formula (Ib) as claimed in claim 13, having one or other of the following characteristics, taken separately or in combination:

- n represents 2 or 3;

- ▶ B represents a phenylene;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

15. The compound of formula (Ib) as claimed in claim 13 or 14, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

16. The compound of formula (Ib) as claimed in any one of claims 13 to 15 simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

17. The compound of formula (Ib) as claimed in any one of claims 13 to 16 simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;

- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

18. The compound of formula (Ib) as claimed in any one of claims 13 to 17 simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ C represents $-\text{O}-$;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

19. The compound as claimed in claim 1 and of formula (Ic)



(Ic)

in which

- ▶ n represents 1, 2 or 3;
- ▶ A represents a substituent chosen from $-\text{C(O)}-$, $-\text{C(S)}-$, $-\text{CH}_2-$, $-\text{CHR}^{10}-$, $-\text{CR}^{10}\text{R}^{11}-$, $-\text{C(O)O}-$, $-\text{C(O)S}-$, $-\text{C(S)O}-$, $-\text{C(S)S}-$, $-\text{C(O)NH}-$, $-\text{C(NH)NH}-$ or $-\text{C(S)NH}-$;
- ▶ B represents
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

- a naphthylene;
- a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
- a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a biphenylene;
- or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;

► E and G represent, independently of each other, a substituent chosen from H, OH, OR^{20} , NH_2 , NHR^{20} ;

► R^1 represents a substituent chosen from H, C_{1-6} -alkyl, $C(O)H$, and $C(O)CH_3$;

► R^2 , R^3 , and R^6 represent, independently of each other, a substituent chosen from H, C_{1-6} -alkyl, $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^4 represents a substituent chosen from H, C_{1-6} -alkyl or R^{21} ;

► R^5 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl or R^{22} ;

► R^7 represents a substituent chosen from H, C_{1-6} -alkyl, arabinosyl or R^{23} ;

► R^8 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$ or R^{24} ;

► R^9 represents a substituent chosen from H, C_{1-6} -alkyl, mannose, glycerol or R^{25} ;

► R^{10} , R^{11} , R^{17} and R^{18} represent, independently of each other, a substituent chosen from C_{1-6} -alkyl or F;

► R^{14} , R^{15} , R^{16} and R^{19} represent, independently of each other, a substituent chosen from H or C_{1-6} -alkyl, $-C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} represent, independently of each other, a substituent chosen from $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are

agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

20. The compound of formula (Ic) as claimed in claim 19, having one or other of the following characteristics, taken separately or in combination:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;
- ▶ B represents a phenylene;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

21. The compound of formula (Ic) as claimed in claim 19 or 20, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- ▶ R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

22. The compound of formula (Ic) as claimed in any one of claims 19 to 21, simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH₂-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH₃;
- ▶ R¹ represents H, CH₃ or C(O)CH₃;
- ▶ R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- ▶ R⁴ represents H, C(O)CH₃ or C(O)NH₂;

- R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

23. The compound of formula (Ic) as claimed in any one of claims 19 to 22, simultaneously having the following characteristics:

- n represents 2 or 3;
- A represents $-\text{C}(\text{O})-$ or $-\text{CH}_2-$;
- B represents a phenylene;
- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent $\text{NHC}(\text{O})\text{CH}_3$;
- R^1 represents H, CH_3 or $\text{C}(\text{O})\text{CH}_3$;
- R^1 represents H or CH_3 ;
- R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- R^4 represents H, $\text{C}(\text{O})\text{CH}_3$ or $\text{C}(\text{O})\text{NH}_2$;
- R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

24. The compound as claimed in claim 1 and of formula (Id)



in which

- n represents 1, 2 or 3;
- B represents
- an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a naphthylene;
 - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
 - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

- a biphenylene;
- or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;

► E and G represent, independently of each other, a substituent chosen from H, OH, OR^{20} , NH_2 , NHR^{20} ;

► R^1 represents a substituent chosen from H, C_{1-6} -alkyl, $C(O)H$, and $C(O)CH_3$;

► R^2 , R^3 , and R^6 represent, independently of each other, a substituent chosen from H, C_{1-6} -alkyl, $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^4 represents a substituent chosen from H, C_{1-6} -alkyl or R^{21} ;

► R^5 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl or R^{22} ;

► R^7 represents a substituent chosen from H, C_{1-6} -alkyl, arabinosyl or R^{23} ;

► R^8 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$ or R^{24} ;

► R^9 represents a substituent chosen from H, C_{1-6} -alkyl, mannose, glycerol or R^{25} ;

► R^{10} , R^{11} , R^{17} and R^{18} represent, independently of each other, a substituent chosen from C_{1-6} -alkyl or F;

► R^{14} , R^{15} , R^{16} and R^{19} represent, independently of each other, a substituent chosen from H or C_{1-6} -alkyl, $-C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} represent, independently of each other, a substituent chosen from $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

25. The compound of formula (Id) as claimed in claim 24, having one or other of the following characteristics, taken separately or in combination:

- n represents 2 or 3;

- ▶ B represents a phenylene;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or CH_3 ;
- ▶ R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

5

10 26. The compound of formula (Id) as claimed in claim 24 or 25, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or CH_3 ;
- ▶ R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

15

20 27. The compound of formula (Id) as claimed in any one of claims 24 to 26, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or CH_3 ;
- ▶ R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

25

30 28. The compound of formula (Id) as claimed in any one of claims 24 to 27, simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent NHC(O)CH_3 ;

35

- R¹ represents H or CH₃;
- R², R³, R⁵, R⁶, R⁷ and R⁹ represent H;
- R⁴ represents H, C(O)CH₃ or C(O)NH₂;
- R⁸ represents H, SO₃H, SO₃Li, SO₃Na, SO₃K, SO₃N(C₁₋₈alkyl)₄, fucosyl or methylfucosyl.

5

29. The compound as claimed in claim 1 and of formula (Ie)



(Ie)

10 in which

- n represents 1, 2 or 3;
- B represents
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a naphthylene;
 - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
 - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
 - a biphenylene;
 - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

15

20

these groups possibly being substituted with one or two substituents R¹² and R¹³ chosen, independently of each other, from halogen, CN, C(O)OR¹⁴, C(O)NR¹⁵R¹⁶, CF₃, OCF₃, -NO₂, N₃, OR¹⁴, SR¹⁴, NR¹⁵R¹⁶ and C₁₋₆-alkyl;

25

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;

► E and G represent, independently of each other, a substituent chosen from H, OH, OR²⁰, NH₂, NHR²⁰;

30

► R¹ represents a substituent chosen from H, C₁₋₆-alkyl, C(O)H, and C(O)CH₃;

► R^2 , R^3 , and R^6 represent, independently of each other, a substituent chosen from H, C_{1-6} -alkyl, $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^4 represents a substituent chosen from H, C_{1-6} -alkyl or R^{21} ;

► R^5 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl or R^{22} ;

► R^7 represents a substituent chosen from H, C_{1-6} -alkyl, arabinosyl or R^{23} ;

► R^8 represents a substituent chosen from H, C_{1-6} -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$ or R^{24} ;

► R^9 represents a substituent chosen from H, C_{1-6} -alkyl, mannose, glycerol or R^{25} ;

► R^{10} , R^{11} , R^{17} and R^{18} represent, independently of each other, a substituent chosen from C_{1-6} -alkyl or F;

► R^{14} , R^{15} , R^{16} and R^{19} represent, independently of each other, a substituent chosen from H or C_{1-6} -alkyl, $-C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

► R^{20} , R^{21} , R^{22} , R^{23} , R^{24} and R^{25} represent, independently of each other, a substituent chosen from $C(O)C_{1-6}$ -alkyl, $-C(S)C_{1-6}$ -alkyl, $-C(O)OC_{1-6}$ -alkyl, $-C(O)NH_2$, $-C(S)NH_2$, $-C(NH)NH_2$, $-C(O)NHC_{1-6}$ -alkyl, $-C(S)NHC_{1-6}$ -alkyl or $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

30. The compound of formula (Ie) as claimed in claim 29, having one or other of the following characteristics, taken separately or in combination:

► n represents 2 or 3;

► B represents a phenylene;

► D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;

► E and G represent $NHC(O)CH_3$;

► R^1 represents H or $C(O)CH_3$;

► R^2 , R^3 , R^5 , R^6 , R^7 and R^9 represent H;

► R^4 represents H, $C(O)CH_3$ or $C(O)NH_2$;

► R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$, fucosyl or methylfucosyl.

31. The compound of formula (Ie) as claimed in claim 29 or 30, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

32. The compound of formula (Ie) as claimed in any one of claims 29 to 31, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

33. The compound of formula (Ie) as claimed in any one of claims 29 to 32, simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent NHC(O)CH_3 ;
- ▶ R^1 represents H or C(O)CH_3 ;
- ▶ $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$ and R^9 represent H;
- ▶ R^4 represents H, C(O)CH_3 or C(O)NH_2 ;
- ▶ R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$, fucosyl or methylfucosyl.

34. The compound as claimed in any one of claims 1 to 33, for which

- ▶ B represents
 - a naphthylene;
 - an arylene;
 - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- or

- a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl.

35. The compound as claimed in any one of claims 1 to 34, for which

► B represents

- an arylene;
- or a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl.

36. The compound as claimed in any one of claims 1 to 35, for which

► B represents

- a phenylene;
- or a heterophenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, $C(O)OR^{14}$, $C(O)NR^{15}R^{16}$, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl.

37. The compound as claimed in any one of claims 1 to 33, for which

► B represents a substituent chosen from:

B1		B6		B11		B16	
B2		B7		B12		B17	
B3		B8		B13		B18	
B4		B9		B14		B19	
B5		B10		B15		B20	

in which R^{12} and R^{13} represent two substituents chosen, independently of each other, from halogen, CN, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl.

38. The compound as claimed in claim 37, for which B represents a phenylene B1 that may be substituted with one or two substituents R^{12} and R^{13} chosen, independently of each other, from halogen, CN, CF_3 , OCF_3 , $-NO_2$, N_3 , OR^{14} , SR^{14} , $NR^{15}R^{16}$ and C_{1-6} -alkyl.

39. The compound as claimed in one of the preceding claims, having one of the following characteristics, taken separately or in combination:


- ▶ $n = 2$ or 3 ;
- ▶ A represents $-C(O)-$ or $-CH_2-$;
- ▶ C represents $-O-$;
- ▶ E and G represent $NHC(O)CH_3$;
- ▶ R^1 represents H or $C(O)CH_3$;
- ▶ R^2 , R^3 , R^5 , R^6 , and R^7 represent a hydrogen atom;
- ▶ R^4 represents a substituent chosen from H, $C(O)CH_3$ and $C(O)NH_2$;

- ▶ R^8 represents a substituent chosen from H, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K and $SO_3N(C_{1-8}alkyl)_4$;
- ▶ R^9 represents a hydrogen atom.

5 40. The compound as claimed in one of the preceding claims, having all of the following characteristics:

- ▶ $n = 2$ or 3 ;
- ▶ A represents $-C(O)-$ or $-CH_2-$;
- ▶ C represents $-O-$;
- ▶ E and G represent $NHC(O)CH_3$;
- ▶ R^1 represents H or $C(O)CH_3$;
- ▶ R^2 , R^3 , R^5 , R^6 , and R^7 represent a hydrogen atom;
- ▶ R^4 represents a substituent chosen from H, $C(O)CH_3$ or $C(O)NH_2$;
- ▶ R^8 represents a substituent chosen from H, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO_3H , SO_3Li , SO_3Na , SO_3K or $SO_3N(C_{1-8}alkyl)_4$;
- ▶ R^9 represents a hydrogen atom.

41. The compound as claimed in one of the preceding claims, for which R^8 represents H, SO_3H , SO_3Li , SO_3Na , SO_3K , $SO_3N(C_{1-8}alkyl)_4$ or a substituent of formula:



20 in which

- ▶ R^{26} represents a substituent chosen from H and CH_3 ;
- ▶ R^{27} and R^{28} represent, independently of each other, a substituent chosen from H, $C(O)CH_3$, SO_3H , SO_3Li , SO_3Na , SO_3K and $SO_3N(C_{1-8}alkyl)_4$.




25 42. The compound as claimed in claim 41, for which R^{26} , R^{27} and R^{28} represent a hydrogen atom.

43. The compound as claimed in one of the preceding claims, for which D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 7 to 15 carbon atoms.



30 44. The compound as claimed in one of the preceding claims, for which D represents a hydrocarbon-based chain according to one of the formulae represented below

46. The compound as claimed in one of the preceding claims, for which D represents a linear hydrocarbon-based chain containing 11 carbon atoms, which is saturated, or unsaturated between carbon atoms 4 and 5.

47. The compound as claimed in one of the preceding claims, corresponding to one of the following formulae:







in which, when it is present, M represents a cation chosen from H⁺, Li⁺, Na⁺, K⁺ and (C₁₋₈alkyl)₄N⁺.